



## The Pattern of Psychoactive Substance Use among Undergraduates in the University of Port Harcourt, Rivers State, Nigeria

C. Okechukwu<sup>1\*</sup>, S. A. Abolo<sup>2</sup>, R. N. Ogo<sup>3</sup> and O. U. Okorie<sup>4</sup>

<sup>1</sup>Department of Paediatrics and Child Health, Rivers State University Teaching Hospital, Port Harcourt, Rivers State, Nigeria.

<sup>2</sup>Public Health Department, Rivers State Ministry of Health, Rivers State, Nigeria.

<sup>3</sup>New Pitt Meadows Medical Clinic, 12181 Harris Road, Pitt Meadows, British Columbia, V3Y 2E9, Canada.

<sup>4</sup>Hospital Management Board, Rivers State Ministry of Health, Rivers State, Nigeria.

### Authors' contributions

This work was carried out in collaboration among all authors. All authors designed the study, performed the statistical analysis and wrote the protocol. Author CO wrote the first draft of the manuscript and managed the literature searches. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/INDJ/2021/v15i330154

Editor(s):

(1) Dr. Takashi Ikeno, National Institute of Mental Health, Japan.

Reviewers:

(1) Daniel D. Faleti, Institute of Nursing Research, Nigeria.

(2) Simona Zaami, "Sapienza" University of Rome, Italy.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/66556>

Original Research Article

Received 19 January 2021

Accepted 27 March 2021

Published 03 April 2021

### ABSTRACT

**Aim:** This study set out to determine the extent of use, abuse and dependence of commonly abused psychoactive substances and their correlates among undergraduates in the University of Port Harcourt.

**Methodology:** In a cross-sectional survey, using multi-stage sampling, undergraduates in the Social sciences faculty of the University of Port Harcourt, Nigeria were selected. A semi-structured questionnaire adapted from the WHO student drug survey proforma was used as an instrument for data collection. The results were analysed using EPI-info 2000 statistical package.

**Results:** There were 352 respondents with a mean age of  $24.4 \pm 3.5$  years. There were 203 (57.7%) males and 149 (42.3%) females. A majority of respondents (90.6%) use at least one psychoactive substance while 22.2% of them abuse substances. The male sex predominated

\*Corresponding author: E-mail: [chiomaokechukwu412@gmail.com](mailto:chiomaokechukwu412@gmail.com);

among the substance abusers with a M: F ratio of 2:1. The prevalence of current use for psychoactive substances ranged from 1.3% to 74.9%. Alcohol (74.9%), was the most commonly used psychoactive substances while heroin (1.3%), was the least used. Abuse rates were higher in cocaine, tobacco and cannabis (66.7%, 63.6% and 55.3% respectively) while dependency rates were higher in heroin, cannabis and cocaine (100.0%, 76.2% and 50.0% respectively). The top three reasons given for starting to use psychoactive substances were experimentation (23.1%), group conformity (23.1%) and curiosity (20.5%) while the reasons given for continuing to use/abuse these substances were to feel good (37.2%), relieve stress (20.0%) and avoid withdrawal symptoms (11.5%). The majority (70.5%) of psychoactive substance users started to use substances before enrolment into the university while more than half of the substance users (56.4%) admitted that being in the university did not increase their use of psychoactive substances. Adverse effects on productivity were noted in 60.3% of substance users.

**Conclusion:** The tendency to abuse substances may begin earlier in childhood and adolescent ages, with the male sex, more vulnerable. More efforts at public enlightenment on the detrimental effects of psychoactive substances should be strategically targeted to include the family unit, primary and secondary school children.

*Keywords: Psychoactive; substance abuse; undergraduates; cannabis; caffeine; heroine; alcohol; tobacco; cocaine.*

## 1. INTRODUCTION

Psychoactive substance abuse refers to the habitual use of drugs not needed for therapeutic purposes, to alter one's mood, affect or state of consciousness [1]. It is a maladaptive pattern of drug or substance use that can result in social, legal, occupational or physical consequences [2]. The relative independence and the desire for group conformity experienced by university undergraduates makes them particularly vulnerable to this harmful practice. Hence, the use of illicit drugs and substances have now become commonplace in most tertiary institutions in Nigeria [3].

The World Health Organization defines substance abuse as the harmful or hazardous use of psychoactive substances including alcohol and illicit drugs [4]. Substance dependence is the psychological or physical need to continue taking a substance or drug [4]. Psychological dependence is the intermittent and recurrent craving for the substance of abuse while physical dependence refers to the development of tolerance for the substance or the necessity to take substances to prevent the development of withdrawal symptoms [4]. Tolerance is the need to take progressively higher doses of a particular substance to achieve a given effect. Withdrawal syndrome implies a constellation of signs and symptoms which occur when drug administration in a physically dependent person is abruptly terminated [4]. Some common examples of psychoactive substances include alcohol, cocaine, opium, cannabinoids, amphetamines,

sedative and hypnotics [5]. Anxiolytics and other stimulants like caffeine, hallucinogens, nicotine, volatile solvents and phencyclidine are also classified as psychoactive substances [5]. The deleterious effects of substance abuse on the student include poor academic performance, impaired concentration, truancy, increased expenses, increased risky sexual behaviour, violence, cultist activities and mental health problems [6-9]. Alcohol and tobacco have been documented as leading risk factors of death among the 15-49 years age group [10].

While some undergraduates commenced the use of psychoactive substances before enrolment into the University, others got introduced to the use of these substances while in the university [11]. Several predisposing risk factors to substance use and abuse may be considered [12]. These factors can be grouped broadly into environmental or social factors and personal factors [12,13]. Environmental factors or social factors include availability/ ease of procuring psychoactive substance, peer pressure, curiosity, idleness and unstable family conditions. Personal factors include age, sex, vulnerable personality and low self-esteem [13,14].

Several studies have been done on psychoactive substance abuse in young adults, however, very few studies have attempted to explore the role of the university community (if any) in influencing the undergraduate to use or abuse psychoactive substances [2,12,15,16]. This study, therefore, sought to determine the extent of use, the abuse and dependency rate of commonly used

psychoactive substances, to identify the reasons for substance use or abuse and to explore the influence of the university community on undergraduates as it relates to substance abuse. This will highlight the relevant key areas of focus for public health interventions to curb the menace of psychoactive substance abuse.

## 2. METHODOLOGY

### 2.1 Study Design and Participants

The descriptive cross-sectional study was conducted at the University of Port Harcourt, Choba (UNIPORT). This institution is the biggest and the only federal-owned university in Rivers state. It has 3 main campuses, - Abuja, Delta and Choba, 19 resident hostels, 7 faculties and an estimated population of over forty thousand students as at the time of this research in October 2006. The sample population consisted of registered undergraduate students of Uniport from 2<sup>nd</sup> to 4<sup>th</sup> year. 352 students, identified by their student identification cards, were recruited into this study. A multi-stage sampling method was used to select eligible respondents over eight weeks. In the first stage, a simple random sampling technique was used to select one faculty out of the 7 faculties in UNIPORT (Humanities, Social sciences, Sciences, Education, Engineering, Management Sciences and Health sciences. The Social sciences faculty was chosen by balloting. The social sciences faculty has four departments (Geography and environmental management, political and administrative studies, sociology and economics). In the second stage, a disproportionate stratified sampling technique was used as 60 respondents were allotted to each of the four departments. A departmental list of students was obtained and 20 students from each level (level 200 to 400) recruited into the study.

The required sample size of 360 was calculated with the appropriate formula [17] using the prevalence rate of 33% in a study by Oshikoya and Alli [11].

### 2.2 Data Collection

A structured self-reporting anonymous questionnaire was designed and used to collect information concerning the respondent's knowledge of substance abuse. The 4-page questionnaire which was modified and adapted from the WHO student drug survey proforma,

[18] contained information on biodata, diagnosis of substance abuse, diagnosis of dependence, withdrawal and the effect of the university on substance abuse. The Diagnosis of substance abuse, dependence and withdrawal were made using the DSM-IV- TR Criteria [19] Ten 5<sup>th</sup> year medical students of UNIPORT were trained on data collection and functioned as assistants. The purpose of the study was explained to the respondents and consent form administered. Questionnaires were distributed to the selected respondents.

### 2.3 Pre-testing

The study instrument was pretested among 50 male and female undergraduates of Rivers State University, another tertiary school in Rivers state Nigeria. The questions were subsequently modified in line with the study objectives.

### 2.4 Statistical Analysis

Data were analysed using EPI-info 2000 statistical package and presented using tables.

### 2.5 Ethical Consideration

Approval for the study was obtained from the Ethical Review Committee of the University of Port Harcourt. Written consent was also obtained from study participants. Participation was entirely voluntary and confidentiality was ensured. Codes rather than participants' names were used as personal identifiers.

## 3. RESULTS

Three hundred and fifty-two students returned properly filled questionnaires giving a response rate of 98%. The age, gender and background distribution of the study participants are shown in Table 1. The mean age of the respondents was 24.4 + 3.5 years. There were 203 (57.7%) males and 149 (42.3%) females with a male to female ratio of 1.4: 1. Most (99.5%) of the respondents were Christians. There were 319 (90.6%) substance users and 78 (22%) substance abusers.

The prevalence of current use for psychoactive substances ranged from 1.3% to 74.9%. Alcohol (74.9%), caffeine (55.8%) and tobacco (14.7%) were the most commonly used psychoactive substances while heroin (1.3%), cocaine (1.9%) and cannabis (10.4%) were the least commonly used (Table 1).

Psychoactive substance abuse rates were higher for cocaine, cannabis and tobacco (66.7%, 63.6% and 55.3% respectively) (Table 2), while substance dependency rates were higher for heroin, cannabis and cocaine (100.0%, 76.2% and 50.0% respectively). Adverse effects on productivity were noted in 60.3% of substance users, (Table 3).

The three most common reasons given for starting to abuse psychoactive substances were group conformity (23.1%), experimentation (23.1%) and curiosity (20.5%) (Table 4), while the top three reasons given for continuing to use/abuse substances were to feel good (37.2%), to relieve stress (25.6%) and to avoid withdrawal symptoms (11.5%), (Table 5).

**Table 1. Demographics and substance use**

	n(%)	Total (n=352)
<b>Age (years)</b>		
15-19	40 (11.4)	
20-24	210 (59.7)	
25-29	96 (27.3)	
30-34	4 (1.1)	
35-39	2 (0.6)	352 (100.0)
<b>Sex</b>		
Male	203 (57.7)	
Female	149(42.3)	352 (100.0)
<b>Background</b>		
Rural	73 (20.7)	
Urban	279 (79.3)	352 (100.0)
<b>Substance use</b>		
Users	319 (90.6)	
Non-users	33 (9.4)	352 (100.0)
<b>Substances being used (n=319)</b>		
Alcohol	239 (74.9)	
Cannabis	33 (10.4)	
Cocaine	6 (1.9)	
Tobacco	47(14.7)	
Heroin	4 (1.3)	
Caffeine	178 (55.8)	
<i>(Multiple responses apply)</i>		

**Table 2. Psychoactive substance abuse rate per substance**

Substance	Number of users	Number of Abusers	Abuse rate (%)
Alcohol	239	62	25.9
Cannabis	33	21	63.6
Cocaine	6	4	66.7
Tobacco	47	26	55.3
Heroin	4	2	50.0
Caffeine	178	35	19.7

**Table 3. Psychoactive substance dependency rate per substance**

Substance	Number of Abusers	Dependants	Dependency rate (%)
Alcohol	62	36	58.1
Cannabis	21	16	76.2
Cocaine	4	2	50.0
Tobacco	26	17	65.4
Heroin	2	2	100.0
Caffeine	35	21	60.0

Table 6 shows the characteristics of substance abusers and the effect of the University on psychoactive substance use. Majority of the psychoactive substance abusers were males (69.2%) and from a rural background (69.2%). Over two-thirds of the respondents who abused psychoactive substances initiated use before enrolment into the university while over half (56.4%) of the substance abusers did not significantly increase their use of psychoactive substances while in the university, (Table 6).

#### 4. DISCUSSION

This study aimed to determine the extent of use of psychoactive substances, the reasons for substance use or abuse and the influence of the university community on undergraduates as it relates to substance abuse. In this study, there was a high prevalence of substance use with over 90% of the undergraduate students using substances. This is higher than the prevalence of

**Table 4. Reasons for starting to use psychoactive substances**

Reasons for starting	Frequency	Percentage
Group conformity	18	23.1
Experimentation	18	23.1
Curiosity	16	20.5
Improve academic performance	14	18.0
Emotional stress	9	11.5
Academic stress	5	6.4
Others	9	11.5
Non- response	7	9.0

*\*Multiple responses apply*

**Table 5. Reasons for continuing to use/abuse psychoactive substances**

Reasons for starting	Frequency	Percentage
To feel good	29	37.2
To relieve stress	20	25.6
To avoid withdrawal symptoms	9	11.5
To concentrate	6	7.7
To function efficiently	4	5.1
Peer pressure/conformity	3	3.9
Others	1	1.3
Non- response	16	20.5

*\*Multiple responses apply*

**Table 6. Characteristics of Substance abusers and the effect of enrolment in the University on psychoactive substance use**

	n(%)	Total (n=78)
<b>Sex</b>		
Male	54 (69.2)	78 (100.0)
Female	24(30.8)	
<b>Background</b>		
Rural	54 (69.2)	78 (100.0)
Urban	24 (30.8)	
<b>Time of onset of substance use</b>		
Before enrolment in the university	55 (70.5)	78 (100.0)
After enrolment in the university	19 (24.4)	
Non-response	4 (5.1)	
<b>Effect of enrolment in the University on substance use</b>		
Increase in the use of psychoactive substances	29 (37.2)	78 (100.0)
No increase in the use of psychoactive substances	44 (56.4)	
Non-response	5(6.4)	

65% obtained in a similar study done by Chukwujekwu [3] in a private tertiary institution in Rivers State, Nigeria. The higher prevalence obtained in this present study may be because the students were selected from the social sciences faculty only unlike in the study by Chukwujekwu [3] in Madonna University where the respondents cut across all the faculties in the university. Moreso, a much lower prevalence of 56% of lifetime use of psychoactive substance was obtained in a study done among medical students in Enugu, Nigeria [20]. The courses offered in the social sciences faculty (eg. Geography, Political science, sociology) are generally considered less tedious than the core sciences (eg. Engineering, Medicine etc), and as such students in social sciences faculty may have more time and energy to engage in social activities within the campus. This avenue for increased peer interaction may result in more peer influence and heightened likelihood to engage in risky behaviours.

In this present study, males predominate among psychoactive substance abusers in a M: F ratio of 2.3:1. This is consistent with previous research findings that the male gender is a risk factor for substance abuse in students [3,21-23]. A possible explanation for this is that the male sex is more given to experimentation and adventurous behaviours. Males tend to take more risks and engage in social groups and activities within the university setting where alcohol and other psychoactive substance use is the norm.

The most commonly used substance in this present study was alcohol. This is in sync with the findings of other researchers who reported high prevalences of alcohol use both in secondary school children and university undergraduates [6,24-27]. Alcohol is often available for purchase in the shops and has a wide price range, allowing for even undergraduate students to afford particular brands. Alcohol is also served regularly in social gatherings attended by these students within the university campus as well as during cult meetings [6,28]. Alcohol has been termed a gateway drug as it often, unfortunately, serves as the beginning point for an adventure into full-blown substance use, abuse, addiction and dependence [3].

The other psychoactive substances commonly used as revealed in this study (Caffeine, Tobacco, Cannabis, Heroin, Cocaine), have also

been reported by previous authors as commonly used by students [5,23,29]. Caffeine, which through its central nervous system stimulant activity, has the potential to increase wakefulness, has become commonly used by students to enable them to study for prolonged hours at night [5]. These students are often unaware of the possible harmful effects of habitual use of caffeine. Tobacco, cannabis, heroin and cocaine, on the other hand, are mainly used to achieve a 'high' feeling where the student becomes temporarily oblivious to any current negative situations [5]. This euphoric effect is desired especially by students who are experiencing some form of frustrations emanating from their family, academics, finances or social life. This present study revealed that substance dependency rates were higher for heroin, cannabis and cocaine. With dependency, the students now require these psychoactive substances to function normally. This takes a toll on the student's finances as these particular substances are expensive. There is then a ripple effect on every aspect of the students' life resulting in stealing, lying, truancy, deteriorating health, poor academic performance, school dropout and even death [30].

The most common reasons given for starting and continuing to abuse substance were out of experimentation, group conformity and curiosity. Other reasons given were to improve academic performance, to cope with academic and emotional stress. These reasons can be classified using a four-factor model of motivation for substance use: "social," "coping," "enhancement," and "conformity" [30]. The findings of this study is comparable to that of other researchers who reported curiosity, experimentation and peer influence as major factors leading to the use and abuse of psychoactive drugs amongst students [5,6,28,29]. In this present study 18% of respondents started to and continued to abuse substance believing it would improve their academic performance. Thirty-five percent of secondary school children in a study in southwest Nigeria, [30] believed using substance would improve their academic performance.

Many studies investigating the use of psychoactive substances in secondary schools have shown an alarming prevalence of use of these substances in secondary schools [5,6,21,23] This further supports the findings in this research, as well as that of Umukoro et al, [31] that many students had been initiated into

the use of these dangerous substances before commencing their tertiary education. Furthermore, more than half of the substance abusers reported that being in the university did not increase their substance abuse habit. This suggests that the university environment did not impact on their decision to continue to use the substances or not. This no table finding is similar to the report of Adeyemo et al. [32] in Benin City, Nigeria, where 79.9% of the undergraduates using psychoactive substances, confessed that the university environment did not influence their drug abuse habit. This is important to note as it further buttresses the fact that much of the negative influence which results in university students using or abusing psychoactive substances perhaps exists outside the four walls of the university. This information is key for the structuring of intervention programs to reduce psychoactive substance abuse. More focus should hence be given to enlightening the family unit, primary and secondary education institutions and the community at large while also sensitizing the university undergraduates.

## 5. CONCLUSION

There is an alarmingly high prevalence of psychoactive substance use and abuse among university students with a large proportion of them initiating use before enrolment into the university. More concerted efforts should be made by the government of Nigeria and all members of the society, at all levels to curb this problem. Enlightenment programs should target children while still in their early ages within their families, primary and secondary educational institutions.

## 6. RECOMMENDATIONS

Programs aimed at curbing the menace of substance abuse should begin at a level before the onset of tertiary education in youths. These programs should especially address youths at an earlier age when their values and habits are still being formed. Youth-friendly centres should be actively functioning and well-funded. Rehabilitation centres should also be set up to help abusers and substance-dependent students amend their lifestyle positively.

## 7. STUDY LIMITATIONS

There was no data triangulation. Information obtained from respondents concerning their use or non-use of substance could not be verified

through any other means. Hence the validity of the information given relies solely on the credibility of the respondents. It was also difficult getting all selected study participants to gather at a location, making data collection tedious.

Also, the research participants were strictly from the Social sciences faculty, hence the findings of this research may not represent the behaviour of all students in UNIPORT.

## 8. LINES OF FUTURE RESEARCH

Further research should inquire about the exact age of initiation of psychoactive substance use and the factors influencing initiation of use in children and young adults.

### CONSENT

It is not applicable.

### ETHICAL APPROVAL

It is not applicable.

### FINANCIAL SUPPORT AND SPONSORSHIP

The authors had no external sources of support. The study was carried out using personal funds.

### ACKNOWLEDGEMENT

The authors appreciate the staff and students of the Social sciences faculty of the University of Port Harcourt for their support during data collection.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Hides L, Dawe S, Kavanagh DJ, Young RM. Psychotic symptom and cannabis release in recent-onset psychosis (prospective study). *Br. J. Psych.* 2006;189: 137-143.
2. McArdle P. Substance use by children and young people. *Arch Dis Child.* 2004;89: 701-4.
3. Chukwujekwu Chidozie Donald. Psychoactive substance use among

- Nigerian students; patterns and sociodemographic correlates. *American Journal of Psychiatry and Neuroscience*. 2017;5(2):22-25.
4. World Health Organization. Substance Abuse; 2014. Available:[http://www.who.int/topics/substance\\_abuse/en/](http://www.who.int/topics/substance_abuse/en/). Accessed December 14th, 2020.
  5. Obiechina GO, Isiguzo BC. Curbing the menace of drug use among secondary school students in Nigeria. *European Journal of Research and Reflection in Educational Sciences*. 2016;4(1):53-63.
  6. Amadi Eric C, Akpelu GO. Effects of drug abuse on the academic performance of secondary school students in Emohua local government area of rivers state. *International Journal of Innovative Healthcare and Research*. 2018;6(1):5-11.
  7. Hingson RW, Zha W, Weitzman ER. Magnitude of and trends in alcohol-related mortality and morbidity among US college students ages 18–24, 1998–2005. *J Stud Alcohol Drugs*. 2009;16:12–20.
  8. Parks K, Collins RL, Derrick JL. The influence of marijuana and alcohol use on condoms behavior: Findings from a sample of young adult female bar drinkers. *Psychol Addict Behav In. Press*; 2012.
  9. Squeglia LM, Pulido C, Wetherill RR, Jacobus J, Brown GG, Tapert SF. Brain response to working memory over three years of adolescence: Influence of initiating heavy drinking. *J Stud Alcohol Drugs*. 2012;73:749–760.
  10. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: A systematic analysis for the global burden of disease study 2010. *Lancet*. 2012;380:2224-2260.
  11. Oshikoya KA, Alli A. Perception of drug abuse among Nigerian undergraduates. *World J. Med. Sci*. 2006;1(2):133-139.
  12. Onigbodun OO, Babalola O. Psychodynamics of psychoactive substance misuse among Nigerian adolescents. *Annals Afri Med*. 2004;3: 111-5.
  13. A study of some causative factors of substance abuse among selected secondary school students in Ibadan Nigeria. *The African Symposium*. 2010; 10(2):4-9.
  14. Taremiyan F, Yaghubi H, Pairavi H, Hosseini SR, Zafar M, Moloodi R. Risk and protective factors for substance use among Iranian university students: A national study *Substance Abuse Treatment, Prevention, and Policy*. 2018; 13(46):1-9.
  15. Eneh AU, Stanley PC. Pattern of substance abuse among secondary school students in Rivers state of Nigeria. *Nig J Med*. 2004;13:36-9.
  16. Egbuonu I, Ezechukwu CC, Chukwuka JO, Uwakwe R. Substance abuse among female senior secondary students in Anambra State, Southeastern Nigeria. *Nig J Clin Pract*. 2004;7:53-5.
  17. Araoye MO. Sample size determination. In: Araoye MO. *Research methodology with statistics for health and social sciences*. Ilorin: Nathadex; 2003.
  18. Smart RG, Hughes PH, Johnston LD, Anumonye A, Khant U, Mora ME, et al. *Methodology for student drug survey: WHO offset publication*. 1980;50.
  19. World Health Organization: The ICD-10 classification of Mental and Behavioral disorder; Diagnostic Criteria for research. World Health Publication, Geneva; 2003.
  20. Ihezue UH. Drug abuse among medical students at a Nigerian University: Part 1. Prevalence and Pattern of use. *Journal of the National Medical Association*. 1988; 80(1):81-85.
  21. Akamnnioo Adoyanto E. Correlates of psychoactive substance use among Nigeria adolescents. *Sahel Medical Journal*. 2015;18(4);192-199.
  22. Champion KE, Teeson M, Newton NC. Patterns and correlates of new psychoactive substance use in a sample of Australian High School students. *Drug and Alcohol Review*. 2016;35:338-344.
  23. Manyike PC, Chinawa JM, Chinawa AT, Herbert AO, Nwokocha ARC, Odutola IO. Correlates for psychoactive substance use among boarding secondary school adolescents in Enugu, South East, Nigeria. *BMC Pediatr*. 2016;16:78.
  24. Kanyoni JM, Gishoma D, Ndahindora V. Prevalence of psychoactive substance use among youth in Rwanda. *BMC research notes*. 2015;8:190. DOI:10.1186/s/3104-015-11482-2
  25. Okoza J, Aluede O. Drug abuse among students of Ambrose Alli University, Ekpoma Nigeria. *Eur J Soc. Sci*. 2009;85-93.



26. Otieno AO, Ofulla A. Drug abuse in Kisumu town Western Kenya Africa J. Food Agr Nutr Dev. 2009;9(3): 846-858.
27. Atwoli L, Mungla PA, Ndungu MN, Kinoti KC, Ogot EM. Prevalence of substance use among college students in Eldoret, Western Kenya. BMC Psychiatry. 2011; 11:1-9.
28. Piwana CN, Haggai MP. Drug abuse and cultism in higher institution of learning. The Case Study of University of Jos; 2007.
29. Duru CB, Oluoha UR, Okafor CC, Diwe KC, Iwu AC, et al. Socio-demographic determinants of psychoactive substance use among students of Tertiary Institutions in Imo State, Nigeria. J Addict Res Ther. 2017;8:345.
30. Idowu A, Aremu AO, Olumide A, Ogunlaja AO. Substance abuse among students in selected secondary schools of an urban community of Oyo-state, South-West Nigeria: Implication for policy action. Afri Health Sci. 2018;18(3):776-785. Available:https://dx.doi.org/10.4314/ahs.v18i3.36
31. Umukoro OL, Taiwo A, Maroh I, Mofoluwake M. Prevalence and patterns of drug abuse among students of Tertiary Institutions in Abeokuta, Ogun State, Nigeria. Int J Psychiatry. 2016;1(1): 1-6.
32. Adeyemo FO, Ohaeri B, Okpala PU, Ogodo O. Prevalence of drug abuse amongst university students in Benin City, Nigeria. Public Health Research. 2016; 6(2):31-37.

© 2021 Okechukwu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle4.com/review-history/66556>